




Nerve Cells and Nerve Impulses

The Cells of the Nervous System

- The human nervous system is comprised of two kinds of cells:
 - Neurons
 - Glia
- The human brain contains approximately 100 billion individual neurons.
- Behavior depends upon the communication between neurons.

Cerebral cortex and associated areas: 12 to 15 billion neurons



Cerebellum: 70 billion neurons

Spinal cord: 1 billion neurons




The Cells of the Nervous System

- Like other cells in the body, neurons contain the following structures:
 - Membrane
 - Nucleus
 - Mitochondria
 - Ribosomes
 - Endoplasmic reticulum


The Cells of the Nervous System

- Neuron cells are similar to other cells of the body but have a distinctive shape.
- A **motor neuron** has its soma in the spinal cord and receives excitation from other neurons and conducts impulses along its axon to a muscle.
- A **sensory neuron** is specialized at one end to be highly sensitive to a particular type of stimulation (touch, temperature, odor etc.)

Dendrite



© 2007 Thomson Higher Education



© 2007 Thomson Higher Education

Fig. 2-6, p. 33

The Cells of the Nervous System

- All neurons have the following major components:
 - Dendrites.
 - Soma/ cell body.
 - Axon.
 - Presynaptic terminals.

The Cells of the Nervous System

- **Dendrites**- branching fibers with a surface lined with synaptic receptors responsible for bringing in information from other neurons.
- Some dendrites also contain **dendritic spines** that further branch out and increase the surface area of the dendrite.

The Cells of the Nervous System

- **Soma** - contains the nucleus, mitochondria, ribosomes, and other structures found in other cells.
 - Also responsible for the metabolic work of the neuron.

The Cells of the Nervous System

- **Axon** - thin fiber of a neuron responsible for transmitting nerve impulses away to other neurons, glands, or muscles.
- Some neurons are covered with an insulating material called the **myelin sheath** with interruptions in the sheath known as **nodes of Ranvier**.

The Cells of the Nervous System

- **Presynaptic terminals** refer to the end points of an axon responsible for releasing chemicals to communicate with other neurons.

The Cells of the Nervous System

- Neurons vary in size, shape, and function.
- The shape of a neuron determines its connection with other neurons and its connections with other neurons.
- The function is closely related to the shape of a neuron.
 - Example: Purkinje cells of the cerebellum branch extremely widely within a single plane

The Cells of the Nervous System



- (Types of glia continued)
 - **Oligodendrocytes & Schwann cells**- build the myelin sheath that surrounds the axon of some neurons.
 - **Radial glia**- guide the migration of neurons and the growth of their axons and dendrites during embryonic development.

The Nerve Impulse

- A **nerve impulse** is the electrical message that is transmitted down the axon of a neuron.
- The impulse does not travel directly down the axon but is regenerated at points along the axon.
- The speed of nerve impulses ranges from approximately 1 m/s to 100 m/s.

The Nerve Impulse

- The **resting potential** of a neuron refers to the state of the neuron prior to the sending of a nerve impulse.
- The membrane of a neuron maintains an **electrical gradient** which is a difference in the electrical charge inside and outside of the cell.



(a)
© 2007 Thomson Higher Education


(b)

The Nerve Impulse

- At rest, the membrane maintains an **electrical polarization** or a difference in the electrical charge of two locations.
 - the inside of the membrane is slightly negative with respect to the outside.
(approximately -70 millivolts)

The Nerve Impulse

- The membrane is **selectively permeable**, allowing some chemicals to pass more freely than others.
- Sodium, potassium, calcium, and chloride pass through channels in the membrane.
- When the membrane is at rest:
 - Sodium channels are closed.
 - Potassium channels are partially closed allowing the slow passage of sodium.




The Nerve Impulse

- The **sodium-potassium pump** is a protein complex that continually pumps three sodium ions out of the cells while drawing two potassium ions into the cell.
 - helps to maintain the electrical gradient.
- The electrical gradient and the concentration gradient work to pull sodium ions into the cell.
- The electrical gradient tends to pull potassium ions into the cells.

Distribution of Ions

Movement of Ions





The Nerve Impulse

- An **action potential** is a rapid depolarization of the neuron.
- Stimulation of the neuron past the threshold of excitation triggers a nerve impulse or action potential.

The Nerve Impulse

- **Voltage-activated channels** are membrane channels whose permeability depends upon the voltage difference across the membrane.
 - Sodium channels are voltage activated channels.
- When sodium channels are opened, positively charged sodium ions rush in and a subsequent nerve impulse occurs.



The Nerve Impulse

- After an action potential occurs, sodium channels are quickly closed.
- The neuron is returned to its resting state by the opening of potassium channels.
 - potassium ions flow out due to the concentration gradient and take with them their positive charge.
- The sodium-potassium pump later restores the original distribution of ions.

The Nerve Impulse

- Local **anesthetic drugs** block sodium channels and therefore prevent action potentials from occurring.
 - Example: Novocain

The Nerve Impulse


- The **all-or-none law** states that the amplitude and velocity of an action potential are independent of the intensity of the stimulus that initiated it.
 - Action potentials are equal in intensity and speed within a given neuron.

The Nerve Impulse


- After an action potential, a neuron has a **refractory period** during which time the neuron resists another action potential.
- The **absolute refractory period** is the first part of the period in which the membrane can not produce an action potential.
- The **relative refractory period** is the second part in which it take a stronger than usual stimulus to trigger an action potential.

The Nerve Impulse

- In a motor neuron, the action potential begins at the **axon hillock** (a swelling where the axon exits the soma).
- **Propagation of the action potential** is the term used to describe the transmission of the action potential down the axon.
 - the action potential does not directly travel down the axon.




(a)



(b)

The Nerve Impulse


- The myelin sheath of axons are interrupted by short unmyelinated sections called **nodes of Ranvier**.
- At each node of Ranvier, the action potential is regenerated by a chain of positively charged ion pushed along by the previous segment.




Cutaway view of axon wrapped in myelin

The Nerve Impulse

- **Saltatory conduction** is the word used to describe this “jumping” of the action potential from node to node.
 - Provides rapid conduction of impulses
 - Conserves energy for the cell
- Multiple sclerosis is disease in which the myelin sheath is destroyed and associated with poor muscle coordination.



(a)



(b)